



cancer.org | 1.800.227.2345

Testicular Cancer Causes, Risk Factors, and Prevention

Learn about the risk factors for testicular cancer and if there are things you can do that might help lower your risk.

Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for testicular cancer.

[q BT 72 720 Td 0 -496.9 Td 0 -26 Td /F1 2i52 5ahs51 0 1 95.35 383 u6 0 0 1 72 529.85 Tanythin](#)

Risk Factors for Testicular Cancer

Different cancers have different risk factors. Some risk factors, like smoking and diet, can be changed. Others, like a person's age or family history, can't be changed.

But having a risk factor, or even many, does not mean that you will get the disease. Just as not having risk factors doesn't mean you won't get the disease. And some people who get the disease may not have had any known risk factors. Even if a person with testicular cancer has a risk factor, it's often very hard to know how much that risk factor contributed to the cancer.

Scientists have found few risk factors that make someone more likely to develop testicular cancer.

- Undescended testicle

- Family history
- HIV infection
- Carcinoma in situ
- Cancer in the other testicle
- Age
- Race and ethnicity

abdomen as opposed to one that has descended at least partway. If cancer does develop, it's usually in the undescended testicle, but about 1 out of 4 cases occur in the normally descended testicle. Because of this, some doctors conclude that cryptorchidism doesn't actually cause testicular cancer, but that there's something else that leads to both testicular cancer and abnormal positioning of one or both testicles.

Orchiopexy may reduce the risk of testicular cancer if it's done when a child is younger, but it's not as clear if it's helpful if the child is older. The best time to do this surgery is not clear. Experts in the United States recommend that orchiopexy be done soon after the child's first birthday for reasons that aren't related to cancer (such as fertility).

Family history

Having a father or brother with testicular cancer increases the risk that you will get it, too. But only a small number of testicular cancers occur in families. Most men with testicular cancer do not have a family history of the disease.

Klinefelter's syndrome is an inherited disease that's also linked to an increased risk of testicular cancer.

HIV infection

Some evidence has shown that men infected with the human immunodeficiency virus (HIV), particularly those with AIDS, are at increased risk. No other infections have been shown to increase testicular cancer risk.

Carcinoma in situ

This condition, described in [What Is Testicular Cancer?](#)¹ often doesn't cause a lump in the testicles or any other [symptoms](#)². It isn't clear how often carcinoma in situ (CIS) in the testicles progresses to cancer. In some cases, CIS is found in men who have a testicular biopsy to evaluate infertility or have a testicle removed because of cryptorchidism. Doctors in Europe are more likely than the doctors in this country to look for CIS. This may be why the numbers for diagnosis and progression of CIS to cancer are lower in the United States than in parts of Europe.

Since we don't know how often CIS becomes true (invasive) cancer, it isn't clear if treating CIS is a good idea. Some experts think that it may be better to wait and see if the disease gets worse or becomes a true cancer. This could allow many men with CIS to avoid the risks and side effects of treatment. When CIS is treated, [radiation](#)³ or

[surgery](#)⁴ (to remove the testicle) is used.

Cancer in the other testicle

A personal history of testicular cancer is another risk factor. About 3% or 4% of men who have been cured of cancer in one testicle will at some point develop cancer in the other testicle.

Age

About half of testicular cancers occur in men between the ages of 20 and 34. But this cancer can affect males of any age, including infants and elderly men.

Race and ethnicity

White, American Indian, and Alaska Native men are several times more likely to get testicular cancer than Black, Asian American, and Pacific Islander men. The reason for this is not known.

Worldwide, the risk of developing this disease is highest among men living in the United States and Europe and lowest among men living in Africa or Asia.

Body size

Several studies have found that tall men have a somewhat higher risk of testicular cancer, but some other studies have not. Most studies have not found a link between testicular cancer and body weight.

Unproven or controversial risk factors

Prior injury or trauma to the testicles and recurrent actions such as horseback riding do not appear to be linked to the development of testicular cancer.

Most studies have not found that strenuous physical activity increases testicular cancer risk. Being [physically active](#)⁵ has been linked with a lower risk of several other forms of cancer as well as a lower risk of many other health problems.

Hyperlinks

1. www.cancer.org/cancer/types/testicular-cancer/about/what-is-testicular-cancer.html
2. www.cancer.org/cancer/types/testicular-cancer/detection-diagnosis-staging/signs-and-symptoms.html
3. www.cancer.org/cancer/types/testicular-cancer/treating/radiation-therapy.html
4. www.cancer.org/cancer/types/testicular-cancer/treating/surgery.html
5. www.cancer.org/cancer/risk-prevention/diet-physical-activity.html

References

American Society of Clinical Oncology. Testicular Cancer: Risk Factors. 09/2016. Accessed at www.cancer.net/cancer-types/testicular-cancer/risk-factors on April 26, 2018.

Boccellino M, Vanacore D, Zappavigna S, et al. Testicular cancer from diagnosis to epigenetic factors. *Oncotarget*. 2017 Sep 18;8(61):104654-104663.

Loveday C, Litchfield K, Levy M, et al. Validation of loci at 2q14.2 and 15q21.3 as risk factors for testicular cancer. *Oncotarget*. 2017;9(16):12630-12638.

National Cancer Institute. Testicular Cancer Screening (PDQ®)—Patient Version. January 30, 2018. Accessed at www.cancer.gov/types/testicular/patient/testicular-screening-pdq#section/_8 on April 26, 2018.

National Comprehensive Cancer Network, Clinical Practice Guidelines in Oncology (NCCN Guidelines®), Testicular Cancer, Version 2.2018 -- February 16, 2018. Accessed at www.nccn.org/professionals/physician_gls/pdf/testicular.pdf on April 26, 2018.

Smith ZL, Werntz RP, Eggener SE. Testicular Cancer: Epidemiology, Diagnosis, and Management. *Mf o e638*.

What Causes Testicular Cancer?

The exact cause of most testicular cancers is not known. But scientists have found that the disease is linked with a number of [other conditions](#). A great deal of research is being done to learn more about the causes.

Researchers are learning how certain changes in a cell's DNA can cause the cell to become cancer. DNA is the chemical in each of our cells that makes up our genes. Genes tell our cells how to function. They are packaged in chromosomes, which are long strands of DNA in each cell. Most cells in the body have 2 sets of 23 chromosomes (one set of chromosomes comes from each parent), but each sperm or egg cell has only 23 chromosomes (one set). When the sperm and egg combine, the resulting embryo has a normal number of chromosomes in each cell, half of which are from each parent. We usually look like our parents because they are the source of our DNA. But DNA affects more than how we look.

Some [genes](#)¹ control when our cells grow, divide into new cells, and die.

- Certain genes that help cells grow and divide are called **oncogenes**.
- Others that slow down cell division or make cells die at the right time are called **tumor suppressor genes**.

Cancers can be caused by changes in chromosomes that turn on oncogenes or turn off tumor suppressor genes.

Most testicular cancer cells have extra copies of a part of chromosome 12 (called *isochromosome 12p* or *i12p*). Some testicular cancers have changes in other chromosomes as well, or even abnormal numbers of chromosomes (often too many). Scientists are studying these DNA and chromosome changes to learn more about which genes are affected and how this might lead to testicular cancer.

Hyperlinks

1. www.cancer.org/cancer/understanding-cancer/genes-and-cancer.html

Last Revised: May 17, 2018

Can Testicular Cancer Be Prevented?

Many men with testicular cancer have no known risk factors. And some of the known risk factors, such as undescended testicles, White race, and a family history of the disease, can't be changed. For these reasons, it's not possible to prevent most cases of this disease at this time.

Experts recommend correcting cryptorchidism in boys for a number of reasons (such as preserving fertility and body image), but it's not clear how much this changes the child's risk for testicular cancer.

Last Revised: May 17, 2018

Written by

The American Cancer Society medical and editorial content team
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345